

MATHEMATICS — HONOURS

Paper : CC-9

(Partial Differential Equation and Multivariate Calculus-II)

Full Marks : 65

*The figures in the margin indicate full marks.*

*Candidates are required to give their answers in their own words  
as far as practicable.*

*All symbols have their usual meaning.*

Group – A

(Marks : 20)

1. Answer all questions with proper justification (one mark for correct answer and one mark for justification) : (1+1)×10

(a) Nature of the partial differential equation (PDE)  $u_{xx}^2 + u_x^2 + \sin u = e^y$  is

(i) non-linear first order

(ii) non-linear second order

(iii) linear first order

(iv) none of these.

(b) Elimination of the arbitrary constants  $a$  and  $b$  from the equation  $\log_e (az - 1) = x + ay + b$  gives the PDE

(i)  $\left(1 + \frac{\partial z}{\partial x}\right) \frac{\partial z}{\partial y} = z \frac{\partial z}{\partial x}$

(ii)  $\left(1 + \frac{\partial z}{\partial y}\right) \frac{\partial z}{\partial x} = x \frac{\partial z}{\partial x}$

(iii)  $\left(1 + \frac{\partial z}{\partial y}\right) \frac{\partial z}{\partial x} = z \frac{\partial z}{\partial y}$

(iv) none of these.

(c) Characteristic curves of the PDE  $u_{xx} + 2u_{xy} + 5u_{yy} + u_x = 0$  is given by

(i)  $y + (1 - 2i)x = c_1, y + (1 + 2i)x = c_2$

(ii)  $y - (1 - 2i)x = c_1, y - (1 + 2i)x = c_2$

(iii)  $y - (1 + 2i)x = c_1, y - (1 - 2i)x = c_2$

(iv) none of these.

(d)  $u_{xx} - \sqrt{x}u_{xy} + xu_{yy} = e^{x/2}$  for all  $x \geq 0$  is

(i) hyperbolic for all values of  $x$ .

(ii) parabolic for all values of  $x$ .

(iii) elliptic for all values of  $x$ .

(iv) parabolic for  $x = 0$  and elliptic for  $x > 0$ .

Please Turn Over

(e)  $(x^2 - y^2 - z^2) \frac{\partial z}{\partial x} + 2xy \frac{\partial z}{\partial y} = 2xz$  has a solution

(i)  $x^2 + y^2 + z^2 = z f\left(\frac{y}{z}\right)$

(ii)  $x^2 - y^2 - z^2 = y f\left(\frac{y}{z}\right)$

(iii)  $x^2 + y^2 + z^2 = f\left(\frac{y}{z}\right)$

(iv)  $x^2 - y^2 - z^2 = z f\left(\frac{y}{z}\right)$

(f) The complete solution of the non-linear partial differential equation  $\left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2 = c^2$  is

(i) a cone

(ii) a cylinder

(iii) a sphere

(iv) none of these.

(g) Value of  $\iint xy \, dx \, dy$  over the region bounded by  $xy = 1$ ,  $y = 0$ ,  $y = x$ ,  $x = 1$  is

(i)  $\frac{1}{8}$

(ii)  $\frac{1}{4}$

(iii) 1

(iv)  $\frac{1}{2}$

(h) If the order of integration  $\int_0^1 dy \int_{x=y}^{x=\sqrt{y}} f(x,y) \, dx$  is interchanged, then it will take the form

(i)  $\int_0^1 dx \int_{y=x^2}^{y=x} f(x,y) \, dx$

(ii)  $\int_0^1 dx \int_x^1 f(x,y) \, dx$

(iii)  $\int_0^1 dx \int_{x^2}^1 f(x,y) \, dx$

(iv) none of these.

(i) If  $\vec{F} = x^2 y \hat{i} + xz \hat{j} + 2yz \hat{k}$ , then the value of  $\text{div}\{\text{curl } \vec{F}\}$  is

(i) 1

(ii) 0

(iii) 2

(iv)  $\hat{i} + \hat{k}$

(j) The work done by a particle in the force field  $\vec{F} = 3x^2 \hat{i} + (2xz - y) \hat{j} + z \hat{k}$  along straight line from  $(0, 0, 0)$  to  $(2, 1, 3)$  is

(i) 16 units

(ii) 22 units

(iii) 14 units

(iv) 42 units.

(3)

X(4th Sm.)-Mathematics-H/CC-9/CBCS

MURALIDHAR GIRLS' COLLEGE  
LIBRARY

Group – B  
(Marks : 21)

Answer *any three* questions.

2. (a) Apply Charpit's method to find the complete integral of the PDE  $(p+q)(px+qy)=1$ .  
(b) Form a PDE by eliminating the arbitrary function  $\phi$  and  $\psi$  from the relation  $u(x, y) = y\phi(x) + x\psi(y)$ .  
4+3
3. Using method of separation of variables solve the PDE  $4z_x + z_y = 3z$  under the condition  $z = 3e^{-y} - e^{-5y}$  at  $x=0$ .  
7

4. Using  $\eta = x + y$  as one of the transformation variable, obtain the canonical form of

$$\frac{\partial^2 u}{\partial x^2} - 2 \frac{\partial^2 u}{\partial x \partial y} + \frac{\partial^2 u}{\partial y^2} = 0$$

and hence solve it.  
7

5. A tightly stretched string of length  $l$  with fixed end points is initially at rest in its equilibrium position, and each of its points is given a velocity  $v$ , which is given by

$$v(x) = \begin{cases} cx, & 0 \leq x < \frac{l}{2} \\ c(l-x), & \frac{l}{2} \leq x \leq l \end{cases}$$

Find the displacement,  $c$  being the wave speed.  
7

6. Solve the following initial boundary value problem

$$u_t = u_{xx} \quad (0 < x < \lambda, t > 0)$$

subject to the conditions  $u(x, 0) = 3 \sin n\pi x$  ( $n$  a +ve integer)

$$u(0, t) = u(\lambda, t) = 0.$$
  
7

Group – C

(Marks : 24)

Answer *any four* questions.

7. Using differentiation under the sign of integration find the value of  $\int_0^{\infty} e^{-a^2 x^2} \cos^2 bx \, dx$ .  
6

8. Evaluate the integral  $\iint \frac{dx \, dy}{(1+x^2+y^2)^2}$  taken over the triangle with vertices at  $(0, 0)$ ,  $(2, 0)$  and  $(1, \sqrt{3})$ .  
6

Please Turn Over

9. Find the value of the integral  $\iiint_E \frac{dx dy dz}{x^2 + y^2 + (z-2)^2}$ , where  $E = \{(x, y, z) : x^2 + y^2 + z^2 \leq 1\}$ . 6
10. Define conservative vector field  $\vec{F}$  and express its relation with the scalar potential  $\phi(x, y, z)$ . Write down Gauss's divergence theorem in case of an irrotational vector over the hemisphere centered at origin. 4+2
11. Find  $\oint_c x dy + y dx$  bounded by the closed contour of astroid with  $x = a \cos^3 t$  and  $y = a \sin^3 t$ . 6
12. Find the surface area of the region common to the intersecting cylinders  $x^2 + y^2 = a^2$  and  $x^2 + z^2 = a^2$ . 6
13. Prove that the volume common to the sphere  $x^2 + y^2 + z^2 = a^2$  and the cylinder  $x^2 + y^2 = ay$  is  $\frac{2}{9}(3\pi - 4)a^3$ . 6