

2021

MATHEMATICS — HONOURS

Seventh Paper

(Module : XIII)

Full Marks : 50

*The figures in the margin indicate full marks.**Candidates are required to give their answers in their own words as far as practicable.* \mathbb{R} , \mathbb{C} respectively denote the set of all real numbers and complex numbers

Group - A

[Analysis - IV]

(Marks : 20)

Answer *any two* questions.

1. (a) Let f and g be positive-valued functions defined on $[a, b]$ such that both f and g have infinite discontinuities only at 'a', both are bounded and R -integrable on $[a + \varepsilon, b]$ for $0 < \varepsilon < b - a$.

If $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = l$, where l is a non-zero real number, prove that $\int_a^b f(x) dx$ and $\int_a^b g(x) dx$ converge or diverge together.

- (b) Test the convergence of $\int_0^1 x^{n-1} \log x dx$. 5+5

2. (a) Examine the convergence of $\int_0^{\pi} \frac{\sin x}{x^p} dx$.

- (b) Show that the integral $\int_0^{\infty} e^{-ax} \frac{\sin x}{x} dx$ is convergent if $a \geq 0$.

- (c) Examine the convergence of $\int_0^1 x^{m-1} (\log_e x)^{n-1} dx$. 3+3+4

Please Turn Over

3. (a) Obtain the Fourier series expansion of $f(x)$ in $[-\pi, \pi]$ where

$$f(x) = \begin{cases} 0, & -\pi \leq x < 0 \\ \frac{1}{4}\pi x, & 0 \leq x \leq \pi \end{cases}$$

Hence show that the sum of the series $1 + \frac{2}{3^2} + \frac{2}{5^2} + \dots$ is $\frac{\pi^2}{8}$.

(b) Evaluate $\int_0^1 dy \int_y^1 e^{x^2} dx$.

7+3

4. (a) Let a function f be defined on a rectangle $R = [0, 1 ; 0, 1]$ as follows :

$$f(x, y) = \begin{cases} \frac{1}{2}, & \text{when } y \text{ is rational} \\ x, & \text{when } y \text{ is irrational} \end{cases}$$

Show that

(i) $\int_0^1 dx \int_0^1 f(x, y) dy$ does not exist, but

(ii) $\int_0^1 dy \int_0^1 f(x, y) dx = \frac{1}{2}$.

- (b) Evaluate $\iint_R \sqrt{|x^2 - 2y|} dx dy$ where $R = [-2, 2 ; 0, 2]$

Or,

Show that $\iint_E \frac{dx dy}{\sqrt{(x+y+1)^2 - 4xy}} = \frac{1}{2} \log_e \left(\frac{16}{e} \right)$

by using the transformation $x = u(1+v)$, $y = v(1+u)$, where E is the triangle with vertices $(0, 0)$, $(2, 0)$, $(2, 2)$. (3+2)+5

Group - B**[Metric Space]****(Marks : 15)**5. Answer *any three* questions :

- (a) (i) Let X be the set of all sequences of real numbers and let $x = \{x_n\}_n$ and $y = \{y_n\}_n$ be any two members of X . Define $d : X \times X \rightarrow \mathbb{R}$ by,

$$d(x, y) = \sum_{n=1}^{\infty} \frac{1}{5^n} \frac{|x_n - y_n|}{1 + |x_n - y_n|}$$

Show that d is a metric on X .

- (ii) Show that in a trivial metric space (X, d) , every subset A of X is both open and closed. 3+2
- (b) (i) Prove that in a metric space (X, d) every open ball is an open set. Is the converse true? Justify your answer.
- (ii) Let X denote the set of all Riemann integrable functions on $[a, b]$. For $f, g \in X$,

$$\text{define } d(f, g) = \int_a^b |f(x) - g(x)| dx. \text{ Is } d \text{ a metric on } X? \text{ Justify.} \quad (2+1)+2$$

- (c) (i) Let (X, d) be a metric space and let A, B be two subsets of X . Then show that

$$\text{diam}(A \cup B) \leq \text{diam}(A) + \text{diam}(B) + d(A, B),$$

where $d(A, B)$ denotes the distance between two sets A and B .

- (ii) Let $A = \{(x, y) : x^2 + y^2 = 2\}$ and $B = \{(x, y) : (x-1)^2 + y^2 = 2\}$.

Find the diameter of the sets $A \cup B$ and $A \cap B$ with respect to the usual metric on \mathbb{R}^2 .

3+2

- (d) (i) Let $\{x_n\}$ be a sequence in a complete metric space (X, d) with the property that

$$\sum_{k=1}^{\infty} d(x_k, x_{k+1}) < \infty. \text{ Show that } \{x_n\} \text{ is convergent.}$$

- (ii) Let $\{x_n\}_n, \{y_n\}_n$ be sequences in a metric space (X, d) such that $x_n \rightarrow x$ in X . Then $y_n \rightarrow x$ in X if and only if $d(x_n, y_n) \rightarrow 0$ in \mathbb{R} .

3+2

- (e) Let (X, d) be a complete metric space and $\{F_n\}_n$ be a descending sequence of non-empty closed

sets in X with diameter of F_n tending to 0 as $n \rightarrow \infty$. Prove that $F = \bigcap_{n=1}^{\infty} F_n$ contains exactly one

point.

5

Please Turn Over

Group - C

[Complex Analysis]

(Marks : 15)

6. Answer *any three* questions :

- (a) (i) Define stereographic projection. Find the image point on the Riemann sphere

$$x^2 + y^2 + \left(z - \frac{1}{2}\right)^2 = \frac{1}{4} \text{ for the point } 4 + 3i \text{ in the complex plane.}$$

- (ii) Show that if
- f
- is analytic in a domain
- $D(\subset \mathbb{C})$
- and
- $|f(z)|$
- is constant in
- D
- , then the function
- $f(z)$
- is constant in
- D
- .
- 2+3

- (b) Let
- G
- be region in
- \mathbb{C}
- and
- $(x_0, y_0) \in G$
- . If
- $u, v : G \rightarrow \mathbb{R}$
- be differentiable at
- (x_0, y_0)
- , prove that,
- $u + iv : G \rightarrow \mathbb{C}$
- is differentiable at
- (x_0, y_0)
- , provided
- u, v
- satisfy the Cauchy–Riemann equations at
- (x_0, y_0)
- .
- 5

- (c) Examine the continuity and differentiability at
- $(0, 0)$
- and the Cauchy–Riemann equations at
- $(0, 0)$
- for the following function defined by

$$f(z) = \begin{cases} \frac{x^5 - y^5}{x^2 + y^2} + i \frac{x^5 + y^5}{x^2 + y^2}, & \text{if } (x, y) \neq (0, 0) \\ 0 & , \text{if } (x, y) = (0, 0) \end{cases} \quad 5$$

- (d) (i) Let
- $u = x^2 - y^2, v = \frac{-x}{x^2 + y^2}$
- . Prove that both
- u
- and
- v
- satisfy Laplace's equation but
- $u + iv$
- is not an analytic function on the complex plane.

- (ii) If
- $f(z)$
- and
- $g(z)$
- be both analytic in a domain
- $D(\subset \mathbb{C})$
- and if
- $f(z)g(z) = U(x, y) + iV(x, y)$
- , then show that
- U
- and
- V
- are both harmonic in
- D
- .
- 2+3

- (e) Define a harmonic function in a region
- $G(\subseteq \mathbb{C})$
- . Prove that
- $u(x, y) = x^3 - 3xy^2$
- is a harmonic function on
- \mathbb{C}
- . Determine its conjugate harmonic and corresponding analytic function.
- 1+2+2