

2023

## MATHEMATICS — HONOURS

Paper : CC-14

(Numerical Methods)

Full Marks : 50

*The figures in the margin indicate full marks.**Candidates are required to give their answers in their own words as far as practicable.*

## Unit - 1

1. Each of the following questions has four possible answers of which exactly one is correct. Choose the correct alternative : 1×10

(a) The maximum absolute error that occurs in rounding off a number to  $m$  decimal places is

(i)  $5 \times 10^{m-1}$

(ii)  $\frac{1}{2} \times 10^{-m-1}$

(iii)  $5 \times 10^{-m-1}$

(iv)  $\frac{1}{2} \times 10^{-m+1}$ .

(b) Find the polynomial of degree  $\leq 3$  passing through the points  $(-1, 1)$ ,  $(0, 1)$ ,  $(1, 1)$  and  $(2, -3)$ .

(i)  $\frac{1}{3}(-2x^3 + 2x + 3)$

(ii)  $-2x^3 + 2x^2 + 2x + 3$

(iii)  $-2x^3 + 2x + 3$

(iv)  $\frac{1}{3}(2x^2 + 2x + 3)$ .

(c) The value of  $\left(\frac{\Delta^2}{E}\right)x^2$  at  $h = 1$  is

(i) 3

(ii) 2

(iii) 1

(iv) 6.

(d) In the Stirling's interpolation formula, the starting point is so chosen that the value of  $u$ , where

$$u = \frac{x - x_0}{h}, \text{ lies between}$$

(i)  $\frac{1}{4} < u < \frac{3}{4}$

(ii)  $\frac{1}{4} < u < 1$

(iii)  $-\frac{1}{4} < u < \frac{1}{4}$

(iv)  $-\frac{3}{4} < u < -\frac{1}{4}$ .

MURALIDHAR GIRLS' COLLEGE  
LIBRARY

Please Turn Over

- (e) Up to which order of polynomial, Simpson's  $\frac{1}{3}$ rd rule provide accurate result?
- (i) 2 (ii) 3  
(iii) 1 (iv) None of these.
- (f) To find the smallest root of the equation  $x^3 = 1 - x^2$  on the interval  $[0, 1]$  by iterative method, the equation should be rewritten as
- (i)  $x = \sqrt{1 - x^3}$  (ii)  $x = \sqrt[3]{1 - x^2}$   
(iii)  $x = \frac{1}{\sqrt{x+1}}$  (iv)  $x = \frac{1 - x^2}{x^2}$ .

- (g) The Runge-Kutta method of order four is used to solve the differential equation  $\frac{dy}{dx} = f(x), y(0) = 0$  with step length  $h$ . The solution at  $x = h$  is given by

(i)  $y(h) = \frac{h}{6} \left[ f(0) + 4f\left(\frac{h}{2}\right) + f(h) \right]$  (ii)  $y(h) = \frac{h}{6} \left[ f(0) + 2f\left(\frac{h}{2}\right) + f(h) \right]$   
(iii)  $y(h) = \frac{h}{6} [f(0) + f(h)]$  (iv)  $y(h) = \frac{h}{6} \left[ 2f(0) + f\left(\frac{h}{2}\right) + 2f(h) \right]$ .

- (h) Let  $A = \begin{pmatrix} 1 & 1 & 2 \\ 1 & 1 & 3 \\ 2 & 3 & 4 \end{pmatrix}, B = \begin{pmatrix} 2 & 2 & 3 \\ 2 & 2 & 4 \\ 9 & 8 & 7 \end{pmatrix}$ . Consider the following two statements.

(P1) LU decomposition for the matrix  $A$  is possible.

(P2) LU decomposition for the matrix  $B$  is possible.

Which of the following statements is true?

- (i) Both P1 and P2 are true (ii) Only P1 is true  
(iii) Only P2 is true (iv) Neither P1 nor P2 is true.
- (i) Power method is applicable on the matrix

(i)  $A = \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}$

(ii)  $A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$

(iii)  $A = \begin{pmatrix} 1 & 1 \\ 0 & 2 \end{pmatrix}$

(iv)  $A = \begin{pmatrix} 0 & -2 \\ -2 & 0 \end{pmatrix}$ .

(3)

Z(6th Sm.)-Mathematics-H/CC-14/CBCS

(j) Which of the following is closed type quadrature ?

(i)  $\int_{-1}^1 f(x)dx = 2f(0)$

(ii)  $\int_{-1}^1 f(x)dx = \frac{1}{2}[f(0) + f(1)]$

(iii)  $\int_{-1}^1 f(x)dx = \frac{1}{2}[f(-1) + f(1)]$

(iv)  $\int_{-1}^1 f(x)dx = \frac{1}{2}[f(-1) + f(0)]$ .

### Unit - 2

Answer *any one* question.

2. (a) Prove that  $(1 + \Delta)(1 - \nabla) = 1$ .

(b) If  $y = 7x^7 - 3x^3$ , find the percentage error in  $y$  at  $x=1$ , if the error in  $x = 0.005$ .

2+3

3. Derive Newton's divided difference interpolation formula for  $(n + 1)$  arguments.

5

### Unit - 3

Answer *any two* questions.

4. Compute  $\frac{dy}{dx}$  and  $\frac{d^2y}{dx^2}$  at  $x=3.8$  using the following table.

$x$	0	1	2	3	4	5	6
$y$	31.23	32.72	33.97	34.74	35.05	34.91	34.51

5

5. (a) Establish the midpoint rule  $\int_a^b f(x)dx = hf\left(a + \frac{h}{2}\right) + \frac{h^3}{24}f''(\xi), a \leq \xi \leq b, h = \frac{b-a}{2}$ .

(b) Evaluate  $\int_0^{10} \frac{dx}{1+x^2}$  by using Simpson's  $\frac{3}{8}$  rule.

3+2

6. (a) If  $T_1, T_2$  denote the Trapezoidal rule approximations to  $I = \int_a^b f(x)dx$  with 1, 2 sub-intervals

respectively, show that  $(I - T_2) = \frac{1}{3}(T_2 - T_1)$ .

(b) What is the degree of precision for Simpson's Three-Eight rule?

4+1

7. Derive the Weddle's rule from Newton-Cotes formula. Also, mention the degree of precision of this method.

4+1

Please Turn Over

## Unit - 4

Answer *any two* questions.

8. Show that the square root of  $N=AB$  is given by  $\sqrt{N} = \frac{S}{4} + \frac{N}{S}$  where  $S = A + B$ . 5
9. Describe Newton's method for solving a system of equations  $f(x, y) = 0$ ,  $g(x, y) = 0$  in two variables  $x$  and  $y$ . When does the method fail? 4+1
10. (a) The equation  $x^3 - 5x^2 + 4x - 3 = 0$  has one root near  $x = 4$  which is to be computed by the following iterative scheme :  $x_{n+1} = \frac{3 + (k-4)x_n + 5x_n^2 - x_n^3}{k}$  with  $x_0 = 4$  and  $k$  as an integer. Determine the value of  $k$  that gives fastest convergence.
- (b) What is / are the difference(s) between the Regula-Falsi and the Secant method. 3+2
11. Show that if the iteration function of the equation  $f(x) = 0$  is such that  $|g'(x)| \leq k < 1$  for all  $x$  in  $[a, b]$ , then the sequence  $\{x_n\}$  generated by  $x_n = g(x_{n-1})$ ;  $n = 1, 2, 3, \dots$  converges to the real root of  $f(x) = 0$  uniquely for any choice of  $x_0$  in  $[a, b]$ . 5

## Unit - 5

Answer *any two* questions.

12. (a) What do you mean by the partial pivoting in solving of system of  $n$  linear equations in  $n$  unknowns? What are the reasons for such pivoting?
- (b) Compute the total number of arithmetic operations (multiplication / division) in Gaussian algorithm for solving an  $(n \times n)$  system of linear equations. (2+1)+2
13. What is the condition of convergence of Gauss-Seidel method? Is it a necessary and sufficient condition? Compare this method with Gauss's elimination method. 2+1+2
14. Describe the power method to calculate the numerically greatest eigenvalue of a real non-singular square matrix of order  $n$ . How do you find its numerically least eigenvalue? 4+1
15. Solve the following system by Crout's method :

$$x + y + z = 3, \quad 2x - y + 3z = 16, \quad 3x + y - z = -3.$$

5

MURALIDHAR GIRLS' COLLEGE  
LIBRARY

## Unit - 6

Answer *any one* question.

16. Using three successive approximations of Picard's method, obtain approximate solution of the differential equation,  $\frac{dy}{dx} = x^2 + y^2$  satisfying the initial condition  $y(0) = 0$ . 5

(5)

Z(6th Sm.)-Mathematics-H/CC-14/CBCS

17. Using Euler's modified method, solve the following differential equation

$$\frac{dy}{dx} = x^2 + y \text{ with } y(0) = 1$$

for  $x = 0.02$  by taking step length  $h = 0.01$ .

5

MURALIDHAR GIRLS' COLLEGE  
LIBRARY